## For Research Use Only

# TYRO3 Polyclonal antibody

Catalog Number: 28513-1-AP 2 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 28513-1-AP BC049368

Concentration: GeneID (NCBI): 7301

Source: UNIPROT ID: Rabbit Q06418

IgG TYRO3 protein tyrosine kinase

Immunogen Catalog Number: Calculated MW:

AG29540 97 kDa

Observed MW: 140 kDa

Full Name:

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

IHC. IF

Isotype:

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse brain tissue, rat brain tissue

IHC: human colon cancer tissue, mouse brain tissue

**Purification Method:** 

WB 1:500-1:2000 IHC 1:50-1:500

IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

IF/ICC: HCT 116 cells,

# **Background Information**

TYRO3 has a structure includes a cytoplasmic domain and an extracellular ligand binding region. And ligand binding at the cell surface induces dimerization and autophosphorylation of TYRO3 on its intracellular domain that provides docking sites for downstream signaling molecules. TYRO3 is associated with many physiological processes including cell survival, migration and differentiation. It has been reported that TYRO3 can be expressed in various regions of the mammalian central nervous system. 28513-1-AP antibody recognizes the 97 kDa and 140 kDa protein which might be the glycosylated form similar to the papers.((PMID: 8873131, 23354874, 21539887)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Quang Duy Trinh	36230987	Cells	IF
Yandan Zhou	38233691	Eur Arch Otorhinolaryngol	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 28513-1-AP (TYRO3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 28513-1-AP (TYRO3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HCT 116 cells using TYRO3 antibody (28513-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).