

For Research Use Only

NOTCH2 Polyclonal antibody, PBS Only

Catalog Number: 28580-1-PBS

Featured Product



Basic Information

Catalog Number:

28580-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG29413

GenBank Accession Number:

NM_024408

GeneID (NCBI):

4853

UNIPROT ID:

Q04721

Full Name:

Notch homolog 2 (Drosophila)

Calculated MW:

265 kDa

Observed MW:

110 kDa, 265 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, Indirect ELISA

Species Specificity:

human

Background Information

NOTCH2 (Neurogenic locus notch homolog protein 2) belongs to the NOTCH family. In kidney, previous reports demonstrated that Notch1 and Notch2 were activated in mammalian nephrogenesis (PMID: 24526233). Notch2 is expressed in many cell types of most lineages in the hematolymphoid compartment and has specific roles in differentiation and function of various immune cells. Notch2 is required for development of splenic marginal zone B cells and regulates differentiation of dendritic cells (DCs) in the spleen. Notch2 appears to play some specific roles in the intestinal immunity, given that the fate of mast cells and a subset of DCs is regulated by Notch2 in the intestine. Notch2 also has important roles in helper T cell divergence from naïve CD4 T cells and activation of cytotoxic T cells (PMID: 22695918).

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




U-87 MG cells were subjected to SDS PAGE followed by western blot with 28580-1-AP (NOTCH2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 28580-1-PBS in a different storage buffer formulation.




WB result of NOTCH2 antibody (28580-1-AP; 1:4000; room temperature for 1.5 hours) with wild-type and NOTCH2 knockout HEK-293T cells. This data was developed using the same antibody clone with 28580-1-PBS in a different storage buffer formulation.




IP result of anti-NOTCH2 (IP:28580-1-AP, 4ug; Detection:28580-1-AP 1:5000) with U-87 MG cells lysate 1960 ug. This data was developed using the same antibody clone with 28580-1-PBS in a different storage buffer formulation.



IP result of anti-NOTCH2 (IP:28580-1-AP, 4ug; Detection:28580-1-AP 1:5000) with HEK-293 cells lysate 1575 ug. This data was developed using the same antibody clone with 28580-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human stomach tissue slide using 28580-1-AP (NOTCH2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 28580-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using NOTCH2 antibody (28580-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 28580-1-PBS in a different storage buffer formulation.