

For Research Use Only

SLC38A1 Polyclonal antibody

Catalog Number: 28632-1-AP

Featured Product



Basic Information

Catalog Number:

28632-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG30088

GenBank Accession Number:

BC010620

GeneID (NCBI):

81539

UNIPROT ID:

Q9H2H9

Full Name:

solute carrier family 38, member 1

Calculated MW:

487 aa, 54 kDa

Observed MW:

54 kDa, 60-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:14000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, mouse brain tissue, mouse skeletal muscle tissue

IHC: mouse brain tissue,

Background Information

SLC38A1 (Sodium-coupled neutral amino acid transporter 1), is also named as SNAT1, which is a symporter that cotransports short-chain neutral amino acids and sodium ions from the extracellular to the intracellular side of the cell membrane (PMID:20599747, PMID:10891391). Inhibition of SLC38A1 reduced tumor growth, cellular migration, invasion, and induced senescence in melanoma cells (PMID:35565278). SLC38A1 was identified as a positive regulator of mTORC1 in neurons, which promoted ischemic brain damage by mTOR-autophagy system (PMID:31552299). SLC38A1 is mainly expressed in brain and placenta.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28632-1-AP (SLC38A1 antibody) at dilution of 1:7000 incubated at room temperature for 1.5 hours.



WB result of SLC38A1 antibody (28632-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SLC38A1 transfected Jurkat cells.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 28632-1-AP (SLC38A1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 28632-1-AP (SLC38A1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).