For Research Use Only

MDMX/MDM4 Polyclonal antibody

Catalog Number: 28747-1-AP 2 Publications



Basic Information

Catalog Number:

28747-1-AP

BC067299

Concentration:

1000 ug/ml

Source:

Rabbit

Cotspan="2">GeneID (NCBI):

4194

Surce:

UNIPROT ID:

Rabbit

Cotspan="2">Cotspan="2">UNIPROT ID:

Rabbit

Cotspan="2">Cotspan="2">Concein Number:

BC067299

Concentration:

GeneID (NCBI):

4194

Source:

Full Name:

Mdm4 p53 binding protein homolog

Immunogen Catalog Number: (mouse)

AG30456 Calculated MW:

490 aa, 55 kDa Observed MW: 70-76 kDa

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB

Species Specificity:

human
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A2780 cells, MCF-7 cells, MDA-MB-231 cells,

Purification Method:

WB 1:2000-1:12000

protein lysate

IHC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

U2OS cells

IP: MCF-7 cells,

IHC: human ovary tumor tissue, human breast cancer

tissue

Background Information

MDM4 is also named as MDMX. It inhibits the activity of the tumor suppressor p53 by primarily cooperating with the p53 feedback regulator MDM2. MDM4 resides mostly in the cytoplasm, but can be recruited to the nucleus by MDM2. (PMID:16511572). The predicted mass of the protein is 54 kDa, while the observed mass is 70-80 kDa, a difference which stated is probably due to phosphorylation or other posttranslational modification(PMID:9226370, 28097652). It can be ubiquitinated and degraded by MDM2(PMID:16163388). It has 5 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhuqian Wang	39437434	J Med Chem	WB
Shiyi Yu	39329452	Cancer Sci	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28747-1-AP (MDMX antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 28747-1-AP (MDM4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-MDMX (IP:28747-1-AP, 4ug; Detection:28747-1-AP 1:3000) with MCF-7 cells lysate 1470 ug.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 28747-1-AP (MDM4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 28747-1-AP (MDMX antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).