For Research Use Only

## SARS-CoV-2 S protein (126-264 aa) Polyclonal antibody



**Purification Method:** 

Antigen affinity purification

Catalog Number: 28869-1-AP

**Basic Information** 

**Applications** 

Catalog Number:

28869-1-AP

Size: 500 μg/ml Source: Rabbit

Isotype:

GenBank Accession Number:

NC\_045512 GeneID (NCBI): 43740568

SARS-CoV-2 Spike Protein

Calculated MW: 141 kDa

Full Name:

Immunogen Catalog Number:

AG30679

**Tested Applications:** 

**ELISA** 

Species Specificity:

Virus

## **Background Information**

Coronaviruses (CoVs) infect human and animals and cause varieties of diseases, including respiratory, enteric, renal, and neurological diseases. CoV uses its spike protein to recognize ACE2 as its receptors and mediate membrane fusion and virus entry into host cells(PMID: 32221306). Each monomer of trimeric S protein is about 180 kDa, and contains two subunits, S1 and S2,S1 recognizes and binds to host receptors, and subsequent conformational  $changes in S2\ facilitate\ fusion\ between\ the\ viral\ envelope\ and\ the\ host\ cell\ membrane\ (PMID:\ 19198616).\ Although\ membrane\ (PMID:\ 19198616).$ the amino acid sequences of the S-glycoprotein were found to be different between the various HCoV, the structures showed high similarity, but the best 3D structural overlap shared by SARS-CoV and SARS-CoV-2, consistent with the shared ACE2 predicted receptor (PMID: 32522207). The spike protein of CoVs can be a target for vaccine and therapeutic development (PMID: 19198616). 28869-1-AP is specific for spike protein of SARS-COV-2, that antigen region is 126-264aa.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



SARS-CoV-2 Spike Antibody (28869-1-AP) tested by ELISA.SARS-CoV-2 Spike protein was coated onto microtiter plates at 0.15 µg/well and then incubated with a dilution series of SARS-CoV-2 Spike Antibody (28869-1-AP). Bound antibodies were detected with HRP conjugated anti-Rabbit IgG followed by incubation with HRP Substrate and then measuring the resulting absorbance at 450 nm.