For Research Use Only

Phospho-mTOR (Ser2481) Polyclonal antibody



Purification Method:

WB: 1:500-1:1000

Antigen affinity purification

Recommended Dilutions:

Catalog Number: 28879-1-AP

11 Publications

Basic Information

Catalog Number: 28879-1-AP Concentration: 350 μg/ml Source:

Isotype:

Rabbit

289 kDa

Applications

Tested Applications: WB, ELISA

Cited Applications:

Species Specificity:

Mouse

Cited Species: human, mouse, rat GenBank Accession Number:

BC117166 GeneID (NCBI):

UNIPROT ID: P42345 Full Name:

FK506 binding protein 12-rapamycin associated protein 1

Calculated MW: Observed MW: 250-289 kDa

Positive Controls:

WB: Insulin treated NIH/3T3 cells,

Background Information

MTOR, also named FRAP1, FRAP2, and RAPT1, belongs to the PI3/PI4-kinase family. MTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. MTOR is the kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino acids. mTORC2 is also activated by growth factors but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 $promotes the serum-induced formation of stress fibers or F-actin. \,mTOR is phosphorylated at Ser 2448 \,via the PI3$ kinase/Akt signaling pathway and autophosphorylated at Ser2481. mTOR plays a key role in cell growth and homeostasis and may be abnormally regulated in tumors.

Notable Publications

Author	Pubmed ID	Journal	Application
Haili Lang	40353431	Eur J Neurosci	WB
Mengqi Wang	40222540	Int J Biol Macromol	WB
Junli Zhang	39502521	Anal Cell Pathol (Amst)	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Nontreated and insulin-treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 28879-1-AP (Phospho-mTOR (Ser2481) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.