

For Research Use Only

Phospho-STAT5A (Tyr694)/STAT5B (Tyr699) Polyclonal antibody

Catalog Number: 28949-1-AP



Basic Information

Catalog Number:

28949-1-AP

Size:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC027036

GeneID (NCBI):

6776

UNIPROT ID:

P42229

Full Name:

signal transducer and activator of transcription 5A

Calculated MW:

794 aa, 92 kDa

Observed MW:

92-100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

WB : GM-CSF treated TF-1 cells,

Background Information

STAT5A, also named as STAT5, belongs to the transcription factor STAT family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5A is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different GHs. Phosphorylation at Tyr694 is obligatory for Stat5 activation. Activation of STAT5A in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of human STAT5A is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of STAT5A in cells.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated TF-1 and GM-CSF treated TF-1 cells were subjected to SDS PAGE followed by western blot with 28949-1-AP (Phospho-STAT5A (Tyr694)/STAT5B (Tyr699) antibody) at dilution of 1:1000 incubated at 4 °C overnight. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



1x10⁶ GM-CSF treated TF-1 cells were intracellularly stained with 0.13 ug Phospho-STAT5A (Tyr694)/STAT5B (Tyr699) Polyclonal antibody (28949-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.13 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.