For Research Use Only

Phospho-TAK1 (Thr187) Polyclonal antibody

Catalog Number:28958-1-AP

4 Publications




| Basic Information | Catalog Number: GenBank Accession Number: 28958-1-AP BC017715 | | Purification Method: Antigen affinity purification | | | |
|------------------------|--|---|---|----|-------------|--|
| | Size: GenelD (NCBI): | | Recommended Dilutions: WB 1:2000-1:10000 | | | |
| | 400 μg/ml 6885 Source: UNIPROT ID: Rabbit 043318 | | | | | |
| | | | | | | |
| | Isotype: IgG | Full Name: mitogen-activated protein kinase kinase kinase 7 | | | | |
| | | | Calculated MW: 579 aa, 64 kDa | | | |
| | | | Observed MW: | | | |
| Applications | Tested Applications: | | Positive Controls: | | | |
| | WB, ELISA Cited Applications: | WB : Calyculin A treated HEK-293T cells, | | | | |
| | WB | | | | | |
| | Species Specificity: | | | | | |
| | Human Cited Species: | | | | | |
| | human | | | | | |
| Background Information | MAP3K7(Mitogen-activated protein kinase kinase kinase 7) is also named as TAK1 and belongs to the MAP kinase kinase kinase subfamily.It plays an important role in the cascades of cellular responses evoked by changes in the environment.It has been linked to interleukin-1 receptor and tumor necrosis factor receptor signaling (PMID: 16186825).It has 4 isoforms (53-55 kDa; 64-70 kDa and 75-80 kDa)produced by alternative splicing. | | | | | |
| Notable Publications | Author | Pubmed ID | Journal | | Application | |
| | Shuhui Meng | 36309313 | Mol Cell Proteomi | cs | WB | |
| | Cong Xiang | 39256356 | Cell Death Dis | | WB | |
| | Cong Liu | 38459169 | Leukemia | | WB | |
| Storage | Storage: Store at -20°C. Stable for or Storage Buffer: PBS with 0.02% sodium azi Aliquoting is unnecessary f | ide and 50% glycerol p | H 7.3. | | | |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated HEK-293T and Calyculin A treated HEK-293T cells were subjected to SDS PAGE followed by western blot with 28958-1-AP (Phospho-TAK1 (Thr187) antibody) at dilution of 1:5000 incubated $4^{\rm C}$ overnight. The membrane was stripped and reblotted with GAPDH antibody as a loading control.