For Research Use Only

Phospho-TSC2 (Thr1462) Polyclonal antibody



Catalog Number: 29000-1-AP

Basic Information

Catalog Number: 29000-1-AP

Size: 300 µg/ml Source: Rabbit

Isotype: IgG GenBank Accession Number:

BC150300
GeneID (NCBI):
7249
UNIPROT ID:
P49815
Full Name:

tuberous sclerosis 2
Calculated MW:

1807 aa, 201 kDa Observed MW: 200 kDa Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Species Specificity: Human, Mouse Positive Controls:

WB: PDGF treated PC-3 cells, PDGF treated NIH/3T3

cells

Background Information

TSC2, also named as Tuberin, plays an essential role in the cellular energy response pathway. TSC2 forms a physical and functional complex with TSC1 that can inhibit the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. It has been reported that TSC2 can be phosphorylated at Thr1462 by Akt/PKB involved in stimulating cell growth and is activated by insulin. (PMID: 14651849, 12172553)

Storage

Storage:


Store at -20°C.

Storage Buffer:


PBS with 0.02% sodium azide, 50% glycerol pH 7.3 and 0.05% BSA

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated PC-3 and PDGF (HZ-1215) treated PC-3 cells were subjected to SDS PAGE followed by western blot with 29000-1-AP (Phospho-TSC2 (Thr1462) antibody) at dilution of 1:1000 incubated at 37° C for 1 hour.



Non-treated NIH/3T3 and PDGF (HZ-1215) treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 29000-1-AP (Phospho-TSC2 (Thr1462) antibody) at dilution of 1:1000 incubated at $37\,^{\circ}$ for 1 hour.