

For Research Use Only

Phospho-TSC2 (Thr1462) Polyclonal antibody



Catalog Number: 29000-1-AP

1 Publications

Basic Information

Catalog Number: 29000-1-AP	GenBank Accession Number: BC150300	Purification Method: Antigen affinity purification
Size: 300 µg/ml	GeneID (NCBI): 7249	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	UNIPROT ID: P49815	
Isotype: IgG	Full Name: tuberous sclerosis 2	
	Calculated MW: 1807 aa, 201 kDa	
	Observed MW: 200 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : PDGF treated PC-3 cells, PDGF treated NIH/3T3 cells
Cited Applications: WB	
Species Specificity: Human, Mouse	
Cited Species: human	

Background Information

TSC2, also named as Tuberin, plays an essential role in the cellular energy response pathway. TSC2 forms a physical and functional complex with TSC1 that can inhibit the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. It has been reported that TSC2 can be phosphorylated at Thr1462 by Akt/PKB involved in stimulating cell growth and is activated by insulin. (PMID: 14651849, 12172553)

Notable Publications

Author	Pubmed ID	Journal	Application
Yue Zhao	38384328	Research (Wash D C)	WB

Storage

Storage:
Store at -20°C.
Storage Buffer:
PBS with 0.02% sodium azide, 50% glycerol pH 7.3 and 0.05% BSA
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated PC-3 and PDGF (HZ-1215) treated PC-3 cells were subjected to SDS PAGE followed by western blot with 29000-1-AP (Phospho-TSC2 (Thr1462) antibody) at dilution of 1:1000 incubated at 37°C for 1 hour.



Non-treated NIH/3T3 and PDGF (HZ-1215) treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 29000-1-AP (Phospho-TSC2 (Thr1462) antibody) at dilution of 1:1000 incubated at 37°C for 1 hour.