

For Research Use Only

Phospho-PDPK1 (Ser241) Polyclonal antibody



Catalog Number: 29241-1-AP

Basic Information

Catalog Number:

29241-1-AP

Size:

200 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC012103

GeneID (NCBI):

5170

UNIPROT ID:

O15530

Full Name:

3-phosphoinositide dependent protein kinase-1

Calculated MW:

556 aa, 63 kDa

Observed MW:

60-63 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, mouse

Positive Controls:

WB : PDGF treated PC-3 cells, PDGF treated NIH/3T3 cells

Background Information

3-Phosphoinositide dependent protein kinase-1 (PDK1 or PDPK1) is a serine-threonine kinase belonging to AGC kinase family. PDK1 plays a critical role in establishing ACD (Asymmetric cell division) in the epithelium (PMID: 27184845). The kinase activity of PDK1 depends on phosphatidylinositol 3-kinase (PI3K), a key intermediate in signaling pathways including those from growth factor receptors and adhesion molecules. Substrates of PDK1, including AKT and the protein kinase C (PKC) isozymes, regulate a number of essential cell functions (PMID: 20027184). PDPK1 has 5 isoforms produced by alternative splicing of 48-63 kDa and is detected as 60-63 kDa.

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide, 50% glycerol pH 7.3 and 0.05% BSA

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated PC-3 and PDGF (HZ-1215) treated PC-3 cells were subjected to SDS PAGE followed by western blot with 29241-1-AP (Phospho-PDPK1 (Ser241) antibody) at dilution of 1:1000 incubated at 4°C overnight.



Non-treated NIH/3T3 and PDGF (HZ-1215) treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 29241-1-AP (Phospho-PDPK1 (Ser241) antibody) at dilution of 1:2000 incubated at 4°C overnight. The membrane was stripped and re-blotted with GAPDH antibody as loading control.