For Research Use Only

## SLC6A19 Polyclonal antibody

Catalog Number:29372-1-AP 2 Publications




| Basic Information      | Catalog Number:<br>29372-1-AP   | GenBank Accession N<br>BC 148801   | lumber:   | Purification Method:<br>Antigen affinity purification |  |
|------------------------|---|--|---|---|--|
|                        | Concentration:  | GenelD (NCBI):   |   | Recommended Dilutions:                                |  |
|                        | 700 ug/ml<br>Source:  | 340024<br>UNIPROT ID:  |   | WB 1:500-1:2000<br>IHC 1:50-1:500                     |  |
|                        | Rabbit  | Q695T7   |   |   |  |
|                        | Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG31076   | Full Name:<br>solute carrier family 6 (neutral amino<br>acid transporter), member 19 |   |   |  |
|                        |   | Calculated MW:<br>634 aa, 71 kDa   |   |   |  |
|                        |   | Observed MW:<br>71 kDa   |   |   |  |
| Applications           | Tested Applications:<br>WB, IHC, ELISA  |  | Positive Controls:  |   |  |
|                        | Cited Applications:   |  | WB : mouse kidney tissue, HeLa cells, rat kidi<br>IHC : human kidney tissue, human small inte |   |  |
|                        | WB, IHC<br>Species Specificity:<br>human, mouse, rat  | tissue   |   |   |  |
|                        | Cited Species:<br>mouse, rat  |  |   |   |  |
|                        | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0  |  |   |   |  |
| Background Information | SLC6A19, also named as BOAT1, encodes system B(O) transmembrane protein that mediates epithelial resorption of<br>neutral amino acids across the apical membrane in kidney and intestine (PubMed: 15044460). The protein contains<br>634 amino acids and has 12 predicted transmembrane regions with highly expression in human kidney and small<br>intestine and minimal in pancreas (PubMed: 15286787). The expression of SLC6A19 in intestinal cells depends on<br>angiotensin converting enzyme 2 which is the cellular receptor for SARS-CoV-2 (PubMed: 32132184). Mutations in<br>SLC6A19 cause Hartnup disease (PMID: 15286787, PMID: 15286788). |  |   |   |  |
| Notable Publications   | Author  | Pubmed ID Journa   | อไ  | Application   |  |
|                        | Biyang Xu   | 39341364 Am J F  | Pathol  | WB  |  |
|                        | Suyun Xu  | 39019414 J Ethno   | opharmacol  | IHC   |  |
| Storage                | Storage:<br>Store at -20°C. Stable for one yes<br>Storage Buffer:<br>PBS with 0.02% sodium azide a<br>Aliquoting is unnecessary for -2  | nd 50% glycerol pH 7.3.  |   |   |  |


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 29372-1-AP (SLC6A19 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 29372-1-AP (SLC6A19 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 29372-1-AP (SLC6A19 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 29372-1-AP (SLC6A19 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 29372-1-AP (SLC6A19 antibody) at dilution of 1:200 (under 40x lens).