

For Research Use Only

LMO7 Polyclonal antibody

Catalog Number: 29392-1-AP



Basic Information

Catalog Number: 29392-1-AP	GenBank Accession Number: NM_005358	Purification Method: Antigen affinity purification
Size: 600 µg/ml	GeneID (NCBI): 4008	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q8WWI1	
Isotype: IgG	Full Name: LIM domain 7	
Immunogen Catalog Number: AG30016	Calculated MW: 193 kDa Observed MW: 140-160 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A549 cells, HeLa cells, mouse lung tissue

IHC : mouse heart tissue, mouse skeletal muscle tissue

Background Information

LMO7, also known as LOMP, FBX20, belongs to the single Calponin-homology (CH) domain-containing protein family, with a single actin-binding CH domain at the amino terminal end (PMID: 19459066). LMO7 also contains a PDZ domain and a carboxyl-terminal LIM domain, both of which have been described as protein-protein interacting domains (PMID: 15140894). LMO7 encodes three splice variants of calculated molecular weights of 196 kDa (P200), 157 kDa (P150), and 100 kDa (P100) (PMID: 26157009).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 29392-1-AP (LMO7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 29392-1-AP (LMO7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 29392-1-AP (LMO7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 29392-1-AP (LMO7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 29392-1-AP (LMO7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).