For Research Use Only

Phospho-PERK/EIF2AK3 (Ser719) Polyclonal antibody

www.ptglab.com

Catalog Number: 29546-1-AP 26 Publications

Basic Information

Catalog Number: 29546-1-AP Concentration:

700 ug/ml Source: Rabbit

Full Name: Isotype: eukaryotic translation initiation factor 2-alpha kinase 3

> Calculated MW: 1116 aa, 125 kDa Observed MW: 140 kDa

BC126354

GeneID (NCBI):

UNIPROT ID:

Q9NZJ5

GenBank Accession Number:

Purification Method:

Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

Applications

Tested Applications: WB, ELISA

Cited Applications:

Species Specificity:

human Cited Species:

human, mouse, rat, zebrafish, bovine

Positive Controls:

WB: λ phosphatase treated HeLa cells, Thapsigargin treated HeLa cells

Background Information

EIF2AK3 encodes the protein kinase RNA-like ER kinase (PERK), a key regulator of the unfolded protein response (UPR) in response to ER stress. Under ER stress conditions, activation of PERK is triggered by the dissociation of glucose-regulated protein (GRP) 78 (also known as BiP) from its luminal domain, followed by oligomerization and autophosphorylation. Phosphorylated PERK subsequently phosphorylates eukaryotic translation initiation factor 2 alpha (eif2 a), to attenuate global protein translation and reduce incoming ER protein load via upregulated ER chaperone expression. (PMID: 35922637, PMID: 32029570)

Notable Publications

Author	Pubmed ID	Journal	Application
Guanjun Li	39901156	Cell Commun Signal	WB
Tiantian Liu	39743026	Free Radic Biol Med	WB
Hongyu Hu	39787915	Bioorg Chem	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated HeLa cells, phosphatase inhibitor treated and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 29546-1-AP (Phospho-PERK/EIF2AK3 (Ser719) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Non-treated HeLa cells and Thapsigargin treated HeLa cells were subjected to SDS PAGE followed by western blot with 29546-1-AP (Phospho-PERK/EIF2AK3 (Ser719) antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.