For Research Use Only

LAG-3/CD223 Polyclonal antibody

IgG

Catalog Number:29548-1-AP 4 Publications



Purification Method:

IHC 1:2000-1:8000

IF-P 1:50-1:500

Antigen affinity purification Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 29548-1-AP NM_002286 Concentration: GeneID (NCBI): 700 ug/ml 3902 Source: **UNIPROT ID:** Rabbit P18627 Isotype: Full Name:

lymphocyte-activation gene 3

Calculated MW: 57 kDa

Applications

Tested Applications: IHC, IF-P, ELISA **Cited Applications:**

Species Specificity:

human **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human tonsillitis tissue. IF-P: human tonsillitis tissue,

Background Information

LAG-3, also known as CD223, is an immune checkpoint molecule that regulates both T-cell activation and $homeostas is.\,LAG-3\ is\ expressed\ on\ activated\ T\ cells,\ NK\ cells,\ regulatory\ T\ cells,\ and\ plasmacytoid\ dendritic\ cells.\ It$ $is \ a \ CD4-related \ molecule \ that \ binds \ MHC \ class \ II. \ LAG-3 \ plays \ an \ important \ role \ in \ modulating \ T \ cell \ expansion \ and \ role \ in \ modulating \ T \ cell \ expansion \ and \ role \$ function, and blockade of LAG-3 with monoclonal antibodies can augment T cell function. (PMID: 15634887; 21086108; 28783703)

Notable Publications

Author	Pubmed ID	Journal	Application
Yuntao Yao	39714755	Cell Oncol (Dordr)	IHC
Yifan Liu	39372371	Clin Transl Immunology	IHC
Jinwu Peng	38115039	Eur J Med Res	IHC

Storage


Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data







Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 29548-1-AP (LAG-3 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using LAG-3/CD223 antibody (29548-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).