For Research Use Only

Phospho-PDH E1 Alpha (Ser232) Polyclonal antibody

www.ptglab.com

Catalog Number:29582-1-AP 6 Publications

Basic Information

Catalog Number: 29582-1-AP

Size: 240 μg/ml Source: Rabbit Isotype:

alpha 1

43 kDa Observed MW: 40 kDa

Applications

Tested Applications: WB, IF/ICC, ELISA Cited Applications:

WB, IF

Species Specificity: Human Cited Species: human, mouse

GenBank Accession Number:

BC002406 GeneID (NCBI): 5160 **UNIPROT ID:** P08559 Full Name:

pyruvate dehydrogenase (lipoamide)

Calculated MW:

Positive Controls:

WB: HeLa cells, Serum-free treated HeLa cells, 10 %

Purification Method:

WB 1:1000-1:8000 IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

fetal bovine serum treated HeLa cells

IF/ICC: \(\lambda\) phosphatase treated HepG2 cells,

Background Information

PDH E1 Alpha (PDHA1), as the major component of PDH, can be phosphorylated and inactivated by PDHK1. PDHA1 contains three serine residues (Ser232, Ser293, or Ser300) that can be reversibly phosphorylated by a dedicated family of four inhibitory pyruvate dehydrogenase kinases (PDHK1-4) and two reactivating phosphatases (PDP1, 2). Hypoxia induces the expression of PDHK1 and PDHK3 and hyperphosphorylates PDHA1. Phosphorylated PDHA1 inactivates the whole pyruvate dehydrogenase complex (PDC), reduces pyruvate entering into the tricarboxylic acid (TCA) cycle for oxidative phosphorylation, enhances the Warburg effect and promotes tumorigenesis. (PMID: 30993888, PMID: 34749809)

Notable Publications

Author	Pubmed ID	Journal	Application
Menglong Wang	35647084	Front Cardiovasc Med	WB
Jie Wang	39550366	Nat Commun	WB
Stephanie Zur Nedden	39332493	Metabolism	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



10 % fetal bovine serum treated HeLa cells and Serum-free treated HeLa cells were subjected to SDS PAGE followed by western blot with 29582-1-AP (Phospho-PDH E1 Alpha (Ser232) antibody) at dilution of 1:4000 incubated at room temperature for 1 hours. The membrane was stripped and reblotted with GAPDH antibody as loading control.



Immunofluorescent analysis of (4% PFA) fixed $^\lambda$ phosphatase treated HepG2 cells using Phospho-PDH E1 Alpha (Ser232) antibody (29582-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).