For Research Use Only

PDIR Polyclonal antibody

Catalog Number: 29593-1-AP

Featured Product

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 29593-1-AP BC001625 GeneID (NCBI): Size: 400 μg/ml 10954 **UNIPROT ID:** Source: Rabbit Q14554

protein disulfide isomerase family A,

Immunogen Catalog Number: AG31479 Calculated MW: 60 kDa Observed MW:

60 kDa

Full Name:

member 5

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

Isotype:

Species Specificity: Human, Mouse, Rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:16000 IHC 1:50-1:500

Purification Method:

Positive Controls:

WB: HeLa cells, LO2 cells, HepG2 cells, mouse liver

tissue, rat liver tissue

IHC: human placenta tissue,

Background Information

 $PDIR (Protein\ disulfide\ isomerase-related\ protein),\ also\ known\ as\ PDIA5\ (Protein\ Disulfide\ Isomerase\ Family\ A$ Member 5), is a member of the PDI gene family and also exhibits chaperone-like activity. PDIR protein is correlated with immune infiltration, which is highly expressed in glioma and participates in glioma progression(PMID: 33664747). PDIR contributes to disulfide bond rearrangement in ATF6 a under stress conditions that necessary for ATF6 a activation upon ER stress(PMID: 24636989).

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Chen	36003400	Front Immunol	WB,IF

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29593-1-AP (PDIR antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 29593-1-AP (PDIR antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 29593-1-AP (PDIR antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.