

For Research Use Only

TNF-alpha Polyclonal antibody

Catalog Number: 29652-1-AP



Basic Information

Catalog Number: 29652-1-AP	GenBank Accession Number: BC107671	Purification Method: Antigen affinity purification
Size: 230 µg/ml	GeneID (NCBI): 24835	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	UNIPROT ID: P16599	
Isotype: IgG	Full Name: tumor necrosis factor (TNF superfamily, member 2)	
Immunogen Catalog Number: AG31037	Observed MW: 26 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : RAW 264.7 cells,
Species Specificity: rat, mouse	

Background Information

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR2. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, ins resistance, and cancer.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



PMA and ionomycin treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 29652-1-AP (TNF-alpha antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.