

For Research Use Only

RPGRIP1L Polyclonal antibody

Catalog Number: 29778-1-AP

3 Publications



Basic Information

Catalog Number:

29778-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG31168

GenBank Accession Number:

NM_015272

GeneID (NCBI):

23322

UNIPROT ID:

Q68CZ1

Full Name:

RPGRIP1-like

Calculated MW:

151 kDa

Observed MW:

151 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IF

Species Specificity:

Human, mouse, rat

Cited Species:

human, zebrafish

Positive Controls:

WB : HeLa cells, mouse testis tissue, HEK-293 cells, Jurkat cells, rat testis tissue

Background Information

RPGRIP1L, also named as FTM, KIAA1005, belongs to the RPGRIP1 family. It negatively regulates signaling through the G-protein coupled thromboxane A2 receptor. RPGRIP1L may be involved in mechanisms like programmed cell death, craniofacial development, patterning of the limbs, and formation of the left-right axis.(PMID:17558409) Defects in RPGRIP1L are the cause of Joubert syndrome type 7 (JBTS7). Defects in RPGRIP1L are the cause of Meckel syndrome type 5 (MKS5).

Notable Publications

Author	Pubmed ID	Journal	Application
Jun Wang	36533556	Dis Model Mech	WB
Manu Ahmed	39282264	bioRxiv	IF
Emma A Hall	36790165	Elife	IF

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29778-1-AP (RPGRIP1L antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 29778-1-AP (RPGRIP1L antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.