

CHD8 Polyclonal antibody

Catalog Number: 29783-1-AP

1 Publications

Basic Information

Catalog Number:

29783-1-AP

Size:

750 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG31425

GenBank Accession Number:

NM_001170629

GeneID (NCBI):

57680

UNIPROT ID:

Q9HCK8

Full Name:

chromodomain helicase DNA binding protein 8

Calculated MW:

290KD

Observed MW:

290 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:12000

IHC 1:250-1:1000

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

Human, Mouse, Rat

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, mouse brain tissue, rat brain tissue

IHC : mouse brain tissue,

Background Information

Chromodomain helicase domain 8 (CHD8) is one of the most frequently mutated and most penetrant genes in the autism spectrum disorder (ASD). CHD8 is located on 14q11.2. It is part of the SNF2H-like ATP-dependent chromatin remodeling enzymes family referred to as CHD (chromodomain helicase DNA binding). CHD8 has two isoforms: CHD8L, a full-length protein of 280 kDa; and CHD8S (Duplin), a 110 kDa protein of the NH2-terminal chromodomain region resulting from alternative splicing (PMID:19151705). CHD8 is essential for development, as homozygote mutant mice die at an embryonic stage. CHD8 is expressed in the mouse at the embryonic stage (E12.5) in different levels in wide regions of the brain (neocortex, forebrain, ventricular, subventricular and mantle zones, rhombic lip (RL), and the isthmus of the cerebellum, as well as in lower RL and floor plate region of the hindbrain, midbrain, diencephalon, hypothalamus, pituitary gland, craniofacial region, and tongue and olfactory epithelium). In the postnatal mouse brain (P20), CHD8 is expressed in the cerebellum, neocortex, hippocampus, hypothalamus, and olfactory bulb (PMID:30277262). Peak expression levels were observed at E18-P7, then gradually decreased to adulthood. Highest expression was found in neurons, and lower levels in astrocyte and astroglia (PMID:30574290). In the mouse, CHD8 is expressed higher in brain compared to other tissues and in the embryo, compared to adult.

Notable Publications

Author	Pubmed ID	Journal	Application
Fenfei Liang	38905100	Cell Rep	WB

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29783-1-AP (CHD8 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 29783-1-AP (CHD8 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 29783-1-AP (CHD8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).