For Research Use Only

Cytokeratin 19 Polyclonal antibody

Catalog Number: 29855-1-AP



Basic Information

Catalog Number: 29855-1-AP

 Size:
 GeneID (NCBI):

 400 ug/ml
 3880

Source: UNIPROT ID:
Rabbit P08727

Isotype: Full Name:
IgG keratin 19

Immunogen Catalog Number: Calculated MW:

AG31469 44KD

Observed MW: 40-44 kDa

NM_002276

GenBank Accession Number:

Purification Method:

IF/ICC 1:200-1:800

Antigen affinity purification Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:10000

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: BxPC-3 cells, HT-29 cells, HepG2 cells, MCF-7

cells, T-47D cells

IHC : human breast cancer tissue, human colon tissue, human thyroid cancer tissue, human normal colon

IF/ICC: HepG2 cells, HeLa cells, A549 cells

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. Keratin 19 is a type I cytokeratin. It is a biochemical marker of skin stem cells in vivo and in vitro.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




BxPC-3 cells were subjected to SDS PAGE followed by western blot with 29855-1-AP (Cytokeratin 19 antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 29855-1-AP (Cytokeratin 19 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 29855-1-AP (Cytokeratin 19 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human normal colon slide using 29855-1-AP (Cytokeratin 19 antibody) at dilution of 1:8000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using Cytokeratin 19 antibody (29855-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).