For Research Use Only

CD8a Polyclonal antibody

Catalog Number: 29896-1-AP 18 Publications



Basic Information

Catalog Number:

29896-1-AP

NM_001081110

Concentration:

600 ug/ml

12525

Source:

Rabbit

P01731

Isotype:

GenBank Accession Number:

NM_001081110

GeneID (NCBI):

12525

UNIPROT ID:

P01731

Full Name:

CD8 antigen, alpha chain

Immunogen Catalog Number: Calculated MW:

AG31942 27KD

Observed MW: 32-35 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:5000 IF-Fro 1:50-1:500

Applications

Tested Applications: WB, IF-Fro, ELISA Cited Applications:

WB

Species Specificity:

mouse
Cited Species:
human, mouse

Positive Controls:

WB: mouse spleen tissue, mouse thymus tissue

IF-Fro: mouse spleen tissue,

Background Information

CD8 is a transmembrane glycoprotein that is predominantly expressed on the surface of cytotoxic T cells, and can also be found on natural killer cells, cortical thymocytes, and dendritic cells. CD8 is composed of two disulfide-linked chains and can be present as a homodimer of CD8 $^{\alpha}$ or as a heterodimer of CD8 $^{\alpha}$ and CD8 $^{\beta}$ (PMID: 3264320; 8253791). The majority of class I-restricted T cells express mostly the CD8 $^{\alpha}$ $^{\beta}$ heterodimer while CD8 $^{\alpha}$ $^{\alpha}$ homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γ $^{\delta}$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Notable Publications

Author	Pubmed ID	Journal	Application
Mengdi Zhang	39898009	Int J Pharm X	
Junhao Wan	39353383	Int Immunopharmacol	
Gaochao Lv	39276489	Bioorg Chem	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29896-1-AP (CD8a antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse spleen tissue using 29896-1-AP (CD8a antibody) at dilution of 1:200 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).