

For Research Use Only

ADAM17 Polyclonal antibody

Catalog Number: 29948-1-AP **1 Publications**



Basic Information

Catalog Number: 29948-1-AP	GenBank Accession Number: BC136783	Purification Method: Antigen affinity purification
Size: 900 µg/ml	GeneID (NCBI): 6868	Recommended Dilutions: WB 1:2000-1:16000
Source: Rabbit	UNIPROT ID: P78536	
Isotype: IgG	Full Name: ADAM metalloproteinase domain 17	
Immunogen Catalog Number: AG32418	Calculated MW: 824 aa, 93 kDa	
	Observed MW: 90-110 kDa	

Applications

Tested Applications: WB, FC, ELISA	Positive Controls: WB: HUVEC cells,
Cited Applications: WB	
Species Specificity: Human, Mouse	
Cited Species: mouse	

Background Information

The ADAMs (A Disintegrin And Metalloprotease) are multidomain transmembrane proteins. One of the first ADAMs implicated in membrane shedding is ADAM-17, which is shown to release the active form of tumor necrosis factor (TNF)-α from its precursor (PMID:18238782). ADAM17 is also named as CSVP, TACE. The full length protein has 9 glycosylation sites, a signal peptide, propeptide and 2 isoforms produced by alternative splicing. The 120-kDa form of ADAM-17 is expressed more frequently and at higher levels in primary breast carcinomas compared with normal breast tissue (PMID:17438092).

Notable Publications

Author	Pubmed ID	Journal	Application
Linfeng Fan	38164152	Theranostics	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29948-1-AP (ADAM17 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



HUVEC cells were subjected to SDS PAGE followed by western blot with 29948-1-AP (ADAM17 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



1X10⁶ HCT 116 cells were intracellularly stained with 0.4 ug Anti-Human ADAM17 (29948-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).