

For Research Use Only

ADAM17 Polyclonal antibody

Catalog Number: 29948-1-AP

4 Publications



Basic Information

Catalog Number:

29948-1-AP

Size:

900 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG32418

GenBank Accession Number:

BC136783

GeneID (NCBI):

6868

UNIPROT ID:

P78536

Full Name:

ADAM metalloproteinase domain 17

Calculated MW:

824 aa, 93 kDa

Observed MW:

90-110 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:16000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human, mouse

Positive Controls:

WB : HUVEC cells, MCF-7 cells, SKOV-3 cells, RAW 264.7 cells

Background Information

The ADAMs (A Disintegrin And Metalloprotease) are multidomain transmembrane proteins. One of the first ADAMs implicated in membrane shedding is ADAM-17, which is shown to release the active form of tumor necrosis factor (TNF)-α from its precursor (PMID:18238782). ADAM17 is also named as CSVP, TACE. The full length protein has 9 glycosylation sites, a signal peptide, propeptide and 2 isoforms produced by alternative splicing. The 120-kDa form of ADAM-17 is expressed more frequently and at higher levels in primary breast carcinomas compared with normal breast tissue (PMID:17438092).

Notable Publications

Author	Pubmed ID	Journal	Application
Shuang Sun	39117070	Arch Biochem Biophys	WB
Li-Mian Song	38948682	Cancer Manag Res	WB
Lindsey E Bazzone	38755212	Nat Commun	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29948-1-AP (ADAM17 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



HUVEC cells were subjected to SDS PAGE followed by western blot with 29948-1-AP (ADAM17 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.