

For Research Use Only

Noggin Polyclonal antibody

Catalog Number: 30023-1-AP

1 Publications



Basic Information

Catalog Number:

30023-1-AP

Size:

600 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG32755

GenBank Accession Number:

BC034027

GeneID (NCBI):

9241

UNIPROT ID:

Q13253

Full Name:

noggin

Calculated MW:

26 kDa

Observed MW:

32 kDa, 64 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

mouse

Positive Controls:

WB : mouse brain tissue, rat brain tissue

Background Information

Noggin is an extracellular polypeptide acting as an antagonist of bone morphogenetic proteins (BMPs) regulating embryonal development. Noggin inhibits activity of BMP-2, -4, -7, -13, and -14. Noggin is present extracellularly in the matrix or retained at the cell surface via interaction with heparin sulfate proteoglycans. In early development stages, Noggin is produced by the Spemann organizer, allowing dorsal-ventral patterning of BMPs (PMID: 8752214). Subsequently, Noggin is expressed by the notochord regulating BMP-4 signaling in neurogenesis. Additionally, Noggin is present during development in the dermal papilla, connective tissue of the hair follicle, lens, retina, and perocular mesenchyme, as well as in the mesoderm lineage regulating development of the bone, cartilage, and muscles.

Notable Publications

Author	Pubmed ID	Journal	Application
Yage Shang	39030543	J Nanobiotechnology	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30023-1-AP (NOG antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



1×10^6 PC-3 cells were intracellularly stained with 0.4 μ g Noggin Polyclonal antibody (30023-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 μ g Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).