

For Research Use Only

CDK10 Polyclonal antibody

Catalog Number: 30061-1-AP



Basic Information

Catalog Number:

30061-1-AP

Size:

750 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG30365

GenBank Accession Number:

BC017342

GeneID (NCBI):

8558

UNIPROT ID:

Q15131

Full Name:

cyclin-dependent kinase 10

Calculated MW:

283 aa, 32 kDa

Observed MW:

38 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

human, mouse

Positive Controls:

WB : A431 cells, K-562 cells, THP-1 cells, U2OS cells, C2C12 cells

IF/ICC : hTERT-RPE1 cells,

Background Information

Cyclin-dependent kinase 10 (CDK10) is a Cdc2-related kinase that was discovered based on its homology to the Cdc2 PSTA1RE amino acid domain (PMID: 8208557). CDK10 plays a pivotal role in the regulation of fundamental cellular processes, including cell proliferation, transcription regulation, and cell cycle regulation. It partners with cyclin M to phosphorylate substrates such as ETS2 and PKN2 in order to modulate cellular growth. Initial reports have indicated that CDK10 may act as a tumor suppressor in breast cancer (PMID: 26392360). Additional studies have demonstrated tumor suppressive and oncogenic roles for CDK10 in malignancies such as hepatobiliary cancers, gastric cancer, glioma, nasopharyngeal carcinoma, and colorectal cancer (PMID: 22326270, PMID: 29512714, PMID: 23740091, PMID: 29845196)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30061-1-AP (CDK10 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed hTERT-RPE1 cells using CDK10 antibody (30061-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CoraLite® 594 ARL13B antibody (CL594-17711, red).