## For Research Use Only

## POGZ Polyclonal antibody

Catalog Number: 30106-1-AP



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 30106-1-AP
 NM\_015100

 Size:
 GeneID (NCBI):

 700 μg/ml
 23126

 Source:
 UNIPROT ID:

Rabbit Q7Z3K3

Isotype: Full Name:
IgG pogo transposable element with ZNF

Immunogen Catalog Number: domain

AG32435 Calculated MW:

155kd Observed MW: 155 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:200-1:800

**Applications** 

Tested Applications: IF/ICC, IHC, IP, WB, ELISA Species Specificity: Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, HepG2 cells, SH-SY5Y cells, mouse

brain tissue IP: HeLa cells,

IHC: mouse brain tissue, mouse testis tissue

IF: HepG2 cells, U-251 cells

## **Background Information**

POGZ, also known as MRD37, is a zinc-finger protein containing 9 C2H2 zinc fingers, a CENP-B (centromere protein-B) domain, and a DDE domain found in the bacterial transposase/retroviral integrase family. POGZ plays a role in mitotic cell cycle progression and is involved in kinetochore assembly and mitotic sister chromatid cohesion. POGZ plays a role in mitotic chromosome segregation by regulating aurora kinase B/AURKB activation and AURKB and CBX5 dissociation from chromosome arms (PMID:20562864). Heterozygous mutation of POGZ causes White-Sutton syndrome (PMID: 26739615). Expect a band approximately 155 kDa in size corresponding to POGZ by western blotting.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30106-1-AP (POGZ antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using POGZ antibody (30106-1-AP) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-POGZ (IP:30106-1-AP, 4ug; Detection:30106-1-AP 1:2000) with HeLa cells lysate 1800 ug.




Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 10 x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed U-251 cells using POGZ antibody (30106-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).