#### For Research Use Only

# BST2 Polyclonal antibody

Catalog Number:30118-1-AP

**Featured Product** 



**Purification Method:** 

WB 1:5000-1:50000 IF-P 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: 30118-1-AP

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 684

 Source:
 UNIPROT ID:

Rabbit Q10589

Isotype: Full Name:
Inc. bone marro

gG bone marrow stromal cell antigen 2

Immunogen Catalog Number:Calculated MW:AG32333180 aa, 20 kDaObserved MW:

30-36 kDa, 50-70 kDa

GenBank Accession Number:

BC033873

**Applications** 

Tested Applications: WB, IF-P, ELISA Species Specificity:

Human, mouse

Positive Controls: WB: HeLa cells,

IF-P: mouse eye tissue,

## **Background Information**

BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. BST2 may play a role in B-cell activation in rheumatoid arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane proteins, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 is 30-36 kDa (PMID: 19196977; 21237475).

## Storage


#### Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


#### **Selected Validation Data**




HeLa cells were subjected to SDS PAGE followed by western blot with 30118-1-AP (BST2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of BST2 antibody (30118-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-BST2 transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using BST2 antibody (30118-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using BST2 antibody (30118-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).