For Research Use Only

RIF1 Polyclonal antibody

Catalog Number:30119-1-AP




Basic Information	Catalog Number: 30119-1-AP	GenBank Accession Numbe NM_001177663.1	r: Purification Method: Antigen affinity purification
	Size: 600 µ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG32422	GeneID (NCBI): 55183 UNIPROT ID: Q5UIPO Full Name: RAP1 interacting factor hon (yeast) Calculated MW: 274 kDa Observed MW: 300 kDa	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:400-1:1600 IF/ICC 1:600-1:2400
Applications	Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity: Human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Alterna retrieval may be performed buffer pH 6.0	WB cell IHC n retrieval with atively, antigen	itive Controls: : A431 cells, HEK-293T cells, HT-29 cells, HeLa s, HepG2 cells, MCF-7 cells : human cervical cancer tissue, mouse testis tissue CC : HeLa cells, HEK-293 cells
Background Information	RIF1 (Rap1-interacting factor 1) is a multifunctional chromosome stability protein implicated in controlling DNA replication and repair. Human RIF1 can interact with protein phosphatase 1 (PP1) and the interaction is crucial for controlling DNA replication by limiting phosphorylation of the MCM complex (PMID: 28077461). RIF1 also acts with PP1 to protect nascent DNA from overdegradation at stalled replication forks (PMID: 31141682). RIF1 promotes non-homologous end joining (NHEI) in double strand break (DSB) repair (PMID: 30443189). Research studies show that RIF1 is overexpressed in breast cancer tissues and is upgraded in NSCLC tissues (PMID: 30237512).		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	I 50% glycerol pH 7.3.	

 For technical support and original validation data for this product please contact:


 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30119-1-AP (RIF1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 30119-1-AP (RIF 1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 30119-1-AP (RIF1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using RIF 1 antibody (30119-1-AP) at dilution of 1:1200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 30119-1-AP (RIF1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).