For Research Use Only

IFN-gamma Polyclonal antibody

Catalog Number: 30293-1-AP 1 Publications



Basic Information

Catalog Number: 30293-1-AP

Source: Rabbit

Isotype:

NM_138880 GeneID (NCBI): 25712 UNIPROT ID:

GenBank Accession Number:

P01581
Full Name:
interferon gamma
Calculated MW:
18 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IHC: 1:50-1:500

FC (Intra): 0.50 ug per 10^6 cells in a

100 µl suspension

Applications

Tested Applications: IHC, FC (Intra), ELISA Species Specificity:

rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

IHC: rat spleen tissue,

FC (Intra): PMA and ionomycin treated rat splenocytes

cells,

Background Information

IFN gamma (IFNG) is a soluble cytokine that is the only member of the type II class of IFN. It is secreted by Th1 cells, cytotoxic T cells and NK cells. The cytokine is associated with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. It plays crucial roles in pathogen clearance. Aberrant IFNG expression is associated with a number of autoinflammatory and autoimmune diseases. It has been identified in many studies as a biomarker for pleural tuberculosis (TB).

Notable Publications

Author	Pubmed ID	Journal	Application
Dwight D Harris	37940431	Surgery	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded rat spleen tissue slide using 30293-1-AP (IFN Gamma antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat spleen tissue slide using 30293-1-AP (IFN Gamma antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 PMA and ionomycin treated rat splenocytes cells were intracellularly stained with 0.5 ug Anti-Rat IFN Gamma (30293-1-AP) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000, and 0.5 ug APC Anti-Rat CD4 (OX-35) (APC-65179, Clone: OX-35), and 0.5 ug rabbit IgG isotype control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.