

For Research Use Only

GPX4 Polyclonal antibody

Catalog Number: 30388-1-AP **25 Publications**



Basic Information

Catalog Number:

30388-1-AP

Size:

550 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG30650

GenBank Accession Number:

BC021567

GeneID (NCBI):

2879

UNIPROT ID:

P36969

Full Name:

glutathione peroxidase 4
(phospholipid hydroperoxidase)

Observed MW:

20-23 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IF 1:50-1:500

Applications

Tested Applications:

WB, IF-P, FC, ELISA

Cited Applications:

WB, IF, FC, IHC

Species Specificity:

Human, Mouse, Rat

Cited Species:

human, rat, mouse, pig

Positive Controls:

WB: HCT 116 cells, HL-60 cells, Jurkat cells, mouse brain tissue, rat brain tissue

IF: rat testis tissue,

Background Information

GPX4 (Phospholipid hydroperoxide glutathione peroxidase, mitochondrial) protects cells against membrane lipid peroxidation and cell death. Required for normal sperm development and male fertility. It has two isoforms about 20KDa and 22KDa, respectively. GPX4 is a monomer, but it has a tendency to form higher mass oligomers (PMID:17630701). It presents primarily in testis.

Notable Publications

Author	Pubmed ID	Journal	Application
Bo Wang	38454322	Mol Med	WB
Xia Xu	38460811	Life Sci	WB
Huan Wang	38432004	Biomaterials	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30388-1-AP (GPX4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 30388-1-AP (GPX4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



1×10^6 HeLa cells were intracellularly stained with 0.4 μ g Anti-Human GPX4 (30388-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 μ g Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat testis tissue using GPX4 antibody (30388-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat testis tissue using GPX4 antibody (30388-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).