

PEX3 Polyclonal antibody

Catalog Number: 30424-1-AP

Basic Information

Catalog Number:

30424-1-AP

Size:

450 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG31516

GenBank Accession Number:

BC014551

GeneID (NCBI):

8504

UNIPROT ID:

P56589

Full Name:

peroxisomal biogenesis factor 3

Calculated MW:

42 kDa

Observed MW:

42 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HEK-293 cells, HepG2 cells, PC-12 cells

IHC: mouse kidney tissue,

Background Information

Peroxisins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. Peroxin 3, also known as PEX3 (peroxisomal biogenesis factor 3) or TRG18, is a 373 amino acid multi-pass membrane protein that anchored in the peroxisomal membrane and essential for peroxisome biogenesis and integrity. Additionally, Peroxin 3 acts as a docking factor for Peroxin 19 and is required for the import of peroxisomal proteins. Defects in the gene encoding Peroxin 3 are the cause of peroxisome biogenesis disorder complementation group 12 (PBD-CG12) and Zellweger syndrome (ZwS), both of which arise from a failure of peroxisomal protein import.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 30424-1-AP (PEX3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 30424-1-AP (PEX3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 30424-1-AP (PEX3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.