## For Research Use Only

## Galectin 3 Polyclonal antibody

Catalog Number: 30450-1-AP

Featured Product



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 30450-1-AP
 BC001120

 Concentration:
 GeneID (NCBI):

 230 ug/ml
 3958

 Source:
 UNIPROT ID:

 Rabbit
 P17931

P17931 Full Name:

gG lectin, galactoside-binding, soluble, 3

Calculated MW: 26 kDa Observed MW: 29-31 kDa

**Applications** 

Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse, rat

Isotype:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: A431 cells, HeLa cells, MCF-7 cells, NIH/3T3 cells,

**Purification Method:** 

WB 1:2000-1:10000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

mouse colon tissue, rat colon tissue

IHC: human thyroid cancer tissue, mouse colon tissue

## **Background Information**

Galectins are a family of animal lectins defined by shared characteristic amino-acid sequences and affinity for  $\beta$  -galactose-containing oligosac-charides (PMID: 8063692). Galectin-3, a member of the  $\beta$ -galactoside-binding proteins, contains one carbohydrate recognition domain (CRD) and a proline- and glycine-rich N-terminal domain through which is able to form oligomers (PMID: 14758078). Galectin-3 is widely expressed in many normal tissues and a variety of tumors. It is found intracellularly in nucleus and cytoplasm or secreted outside of cell, being present on the cell surface or in the extracellular space (PMID: 16478649). Galectin-3 is involved in various biological processes including cell growth, adhesion, differentiation, apoptosis, angiogenesis, immune response, neoplastic transformation and metastasis (PMID: 16478649; 14758078).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 30450-1-AP (Galectin 3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of Galectin 3 antibody (30450-1-AP; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Galectin 3 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 30450-1-AP (Galectin 3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 30450-1-AP (Galectin 3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).