For Research Use Only

## MUC20 Polyclonal antibody

Catalog Number:30457-1-AP





Basic Information	Catalog Number: 30457-1-AP	GenBank Accession Number: BC029267	Purification Method: Antigen affinity purification		
	Size: 600 µ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG31867	GeneID (NCBI): 200958 UNIPROT ID: Q8N307	Recommended Dilutions: WB 1:1000-1:4000 IF/ICC 1:50-1:500		
				Full Name: mucin 20, cell surface associated Calculated MW: 55 kDa, 71 kDa	
		Observed MW: 70 kDa			
		Applications	Tested Applications: IF/ICC, WB, ELISA Species Specificity: Human, mouse	Positive (	Controls:
				WB : mouse kidney tissue, mouse liver tissue IF/ICC : HepG2 cells,	
Background Informatior	MUC20 contains serine, threonine, and proline-rich repeats in its extracellular domain and is likely to be extensively O-glycosylated. MUC20 blocks GRB2 recruitment to MET thus suppressing the GRB2-RAS pathway and inhibiting HGF-induced proliferation of MMP1 and MMP9 expression (PMID: 15314156). MUC20 is highly expressed in the kidney and localized in the proximal tubules but not in the glomerulus or distal tubules(PMID: 14565953). MUC20 was detected in most of the male urogenital tract epithelia, except for epididymis(PMID: 16997931). The expression of MUC20 is up-regulated in the kidneys of patients with immunoglobulin A nephropathy, in an animal model of lupus nephritis, and in mice with acute renal injury caused by cisplatin administration or unilateral ureteral obstruction(PMID: 14565953).				
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	1 50% glycerol pH 7.3.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 30457-1-AP (MUC20 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunofluorescent analysis of (-20°C Ethanol) fixed HepC2 cells using MUC20 antibody (30457-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).