

For Research Use Only

IL-12/IL-23 p40 Polyclonal antibody

Catalog Number: 30537-1-AP



Basic Information

Catalog Number:

30537-1-AP

Concentration:

400 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001303244

GeneID (NCBI):

16160

UNIPROT ID:

P43432

Full Name:

interleukin 12b

Calculated MW:

38kd

Observed MW:

45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

mouse, rat

Positive Controls:

WB : mouse brain tissue, mouse heart tissue, mouse kidney tissue, rat brain tissue, rat heart tissue

Background Information

Interleukin (IL)-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R β 1 and IL-12R β 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R β 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30537-1-AP (IL-12/IL-23 p40 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.