For Research Use Only

## CD200 Polyclonal antibody

Catalog Number:30570-1-AP




| Basic Information                                      | Catalog Number:<br>30570-1-AP  | GenBank Accession Number:<br>BC022522                      | Purification Method:<br>Antigen affinity purification |              |
|--|--|--|---|--------------|
|  | Size:<br>400 µg/ml   | GenelD (NCBI):<br>4345                                     | Recommended Dilutions:<br>WB 1:500-1:1000             |              |
|  | Source:<br>Rabbit  | UNIPROT ID:<br>P41217                                      |   |              |
|  | lsotype:<br>IgG  | Full Name:<br>CD200 molecule                               |   |              |
|  |  | Calculated MW:<br>269 aa, 30 kDa<br>Observed MW:<br>50 kDa |   |              |
|  |  |  |   | Applications |
| WB, ELISA<br>Species Specificity:<br>Human, Mouse, Rat | WB : mouse brain tissue, rat brain tissue  |  |   |              |
| Background Information                                 | CD200, also known as OX2, is a type I transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 3032785). It contains two extracellular immunoglobulin domains, a transmembrane and a cytoplasmic domain. CD200 is expressed on a variety of cell types, including thymocytes, B lymphocytes, a subset of T lymphocytes, follicular dendritic cells, neurons, and endothelial cells (PMID: 3032785; 10981966). CD200 bind to its receptor, CD200R, which is primarily expressed by myeloid and T cell lineage (PMID: 15187158). The CD200-CD200R interaction plays a role in immunosuppression and suppression of anti-tumor immune responses (PMID: 22020332). |  |   |              |
| Storage  | Storage:<br>Store at -20°C. Stable for one year after shipment.<br>Storage Buffer:<br>PBS with 0.02% sodium azide and 50% glycerol pH 7.3.<br>Aliquoting is unnecessary for -20°C storage  |  |   |              |

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30570-1-AP (CD200 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.