For Research Use Only

IL18RAP Polyclonal antibody

Catalog Number: 30587-1-AP



Basic Information

Catalog Number:

30587-1-AP

BC069630

Concentration:

400 ug/ml

Source:

Rabbit

Cop5256

Isotype:

GenBank Accession Number:

GeneID (NCBI):

8807

UNIPROT ID:

O95256

Full Name:

interleukin 18 receptor accessory

Immunogen Catalog Number: protein

AG30327 Calculated MW:

599 aa, 68 kDa Observed MW: 60 kDa

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HL-60 cells, K-562 cells, RAW 264.7 cells, rat

Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

spleen tissue

IHC: mouse spleen tissue, rat spleen tissue

Background Information

Interleukin-18 receptor accessory protein (IL18RAP, also known as IL1R7 and CD218b) is a crucial component of the IL-18 receptor complex. It enhances the binding activity of IL-18 to its receptor and plays a significant role in IL-18 signaling (PMID: 24842757). IL18RAP is related to the human tumor microenvironment and promotes various immune cell infiltration (PMID: 37698530).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30587-1-AP (IL18RAP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 30587-1-AP (IL18RAP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat spleen tissue slide using 30587-1-AP (IL18RAP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).