

## gp130/IL6ST Polyclonal antibody

Catalog Number: 30591-1-AP

## Basic Information

## Catalog Number:

30591-1-AP

## Size:

550 ug/ml

## Source:

Rabbit

## Isotype:

IgG

## GenBank Accession Number:

NM\_002184.3

## GeneID (NCBI):

3572

## UNIPROT ID:

P40189-1

## Full Name:

interleukin 6 signal transducer  
(gp130, oncostatin M receptor)

## Calculated MW:

104 kDa

## Observed MW:

130-150 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:1000

IF/ICC 1:50-1:500

## Applications

## Tested Applications:

WB, IF/ICC, ELISA

## Species Specificity:

human

## Positive Controls:

WB : A549 cells, HepG2 cells, NCCIT cells

IF/ICC : HepG2 cells,

## Background Information

Glycoprotein 130 (gp130, also known as IL6ST, CD130 or IL6-beta) is a ubiquitously expressed, signal-transducing receptor that serves as the signal transduction unit for IL-6 family of cytokines, including IL-6, IL-11, IL-27, leukemia inhibitory factor (LIF), OSM, ciliary neurotrophic factor (CNTF), cardiotrophin 1 (CT-1), and cardiotrophin-like cytokine (CLC) (PMID: 9143707; 20610800). These cytokines signal through the gp130/Jak/STAT pathway (PMID: 9716487). Binding of IL-6 to IL-6R induces gp130 homodimerization and formation of a high-affinity receptor complex, which activates Jaks. That causes phosphorylation of gp130 tyrosine residues which in turn activates STAT3. gp130 is a type I transmembrane protein, and can also exist as a soluble form (sgp130). sgp130 binds to sIL-6R/IL-6 complexes and prevents their interactions with membrane-anchored gp130 on target cells (PMID: 9815830).

## Storage

## Storage:


Store at -20°C. Stable for one year after shipment.

## Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30591-1-AP (gp130/IL6ST antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using gp130/IL6ST antibody (30591-1-AP) at dilution of 1:200 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).