

For Research Use Only

# Fas/CD95 Polyclonal antibody

Catalog Number:30601-1-AP



## Basic Information

**Catalog Number:**

30601-1-AP

**Size:**

450 µg/ml

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG30426

**GenBank Accession Number:**

NM\_007987

**GeneID (NCBI):**

14102

**UNIPROT ID:**

P25446

**Full Name:**

Fas (TNF receptor superfamily member 6)

**Calculated MW:**

37 kDa

**Observed MW:**

30-40 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:1000-1:4000

IF 1:200-1:800

## Applications

**Tested Applications:**

IF/ICC, WB, ELISA

**Species Specificity:**

Human, mouse, rat

**Positive Controls:**

**WB :** Jurkat cells, J774A.1 cells, mouse brain tissue, rat brain tissue

**IF :** Raji cells,

## Background Information

FAS, also named as CD95, APO-1, APT1, FAS1 and TNFRSF6, is a receptor for TNFSF6/FASLG. It is a cell surface receptor belonging to the TNF receptor superfamily, can mediate apoptosis by ligation with an agonistic anti-Fas antibody or Fas ligand. Stimulation of Fas results in the aggregation of its intracellular death domains, leading to the formation of the death-inducing signaling complex (DISC). FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30601-1-AP (Fas/CD95 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 30601-1-AP (Fas/CD95 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed Raji cells using Fas/CD95 antibody (30601-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).