

For Research Use Only

Lipocalin-2/NGAL Polyclonal antibody

Catalog Number: 30700-1-AP

Featured Product

1 Publications



Basic Information

Catalog Number:

30700-1-AP

Concentration:

400 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG33369

GenBank Accession Number:

GeneID (NCBI):

3934

UNIPROT ID:

P80188

Full Name:

Lipocalin 2

Observed MW:

22-25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

IHC

Species Specificity:

human, mouse

Cited Species:

mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HT-29 cells,

IHC : human colon cancer tissue, Human bowens disease

IF/ICC : RAW 264.7 cells,

Background Information

Lipocalin-2 (LCN-2) is a adipocytokine also referred to as neutrophil gelatinase-associated lipocalin (NGAL). Lipocalin-2 is a circulatory protein responsible for the transportation of small and hydrophobic molecules to target organs. Lipocalin-2 is used as a biomarker for acute and chronic renal injury. It is present in a large variety of cells including neutrophil, hepatocytes, lung, bone marrow, adipose tissue, macrophages, thymus, non-neoplastic breast duct, prostate, and renal cells. Different functions have been associated with Lipocalin-2, including antibacterial, anti-inflammatory, and protection against cell and tissue stress (PMID:34463264).

Notable Publications

Author	Pubmed ID	Journal	Application
Mengqiu Lu	39891958	Redox Biol	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30700-1-AP (Lipocalin-2/NGAL antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




WB result of NGAL antibody (30700-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NGAL transfected HT-29 cells.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 30700-1-AP (Lipocalin-2/NGAL antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human brown disease slide using 30700-1-AP (NGAL antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed RAW 264.7 cells using Lipocalin-2/NGAL antibody (30700-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).