

For Research Use Only

Nestin Polyclonal antibody

Catalog Number: 30760-1-AP



Basic Information

Catalog Number: 30760-1-AP	GenBank Accession Number: NM_006617	Purification Method: Antigen affinity purification
Size: 550 ug/ml	GeneID (NCBI): 10763	Recommended Dilutions: WB 1:500-1:3000
Source: Rabbit	UNIPROT ID: P48681	IHC 1:250-1:1000
Isotype: IgG	Full Name: nestin	IF-P 1:50-1:500
Immunogen Catalog Number: AG29600	Calculated MW: 177 kDa	IF/ICC 1:200-1:800
	Observed MW: 240-280 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA	Positive Controls: WB : SGC-7901 cells, SH-SY5Y cells
Species Specificity: human	IHC : rat cerebellum tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IF-P : mouse brain tissue,
	IF/ICC : SH-SY5Y cells,

Background Information

Nestin, also named as NES and Nbla00170, is a class VI intermediate filament protein expressed in stem cells of the central nervous system (CNS) but not in mature CNS cells. Its expression is used extensively as a marker for neural lineage cells derived from human embryonic stem (hES) and induced pluripotent stem (iPS) cells. Nestin is also expressed in non-neural stem cell populations such as pancreatic islet progenitors and hematopoietic progenitors. Nestin can be detected as 240-280 kDa. Studies point toward nestin heavy posttranslational modifications as possibly serving particular functions in podocytes and in a few other particular cell types(PMID: 17210924).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30760-1-AP (Nestin antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue slide using 30760-1-AP (Nestin antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using NES antibody (30760-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using Nestin antibody (30760-1-AP) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).