For Research Use Only

KIF15 Polyclonal antibody

Catalog Number: 30774-1-AP

Featured Product



Basic Information

 Catalog Number:
 GenBank Accession Number:

 30774-1-AP
 NM_020242

 Concentration:
 GeneID (NCBI):

 700 ug/ml
 56992

 Source:
 UNIPROT ID:

 Rabbit
 Q9NS87

 Isotype:
 Full Name:

IgG kinesin family member 15

Immunogen Catalog Number: Calculated MW: AG33897 160 kDa
Observed MW:

145-150 kDa

Applications

Tested Applications: WB, IF/ICC, ELISA Species Specificity:

human

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cels, K-562 cells,

Purification Method:

WB 1:2000-1:16000 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

LO2 cells

IF/ICC: HEK-293 cells,

Background Information

KIF15, also named as KLP2 and KNSL7, belongs to the kinesin-like protein family and KLP2 subfamily. It is a plus-end directed kinesin-like motor enzyme which involved in mitotic spindle assembly. It is located in the cytoplasm and biased expression in testis. Several isoforms of KIF15 exist due to the alternative splicing, with molecular weight between 130-160 kDa. Sometimes higher molecular weight around 200 kDa can also be observed, which may be a modified variant of KIF15 (14618103). It is known that KIF15 participates in important events during neuronal development and involves multiple cellular processes such as proliferation, apoptosis, and differentiation. There is emerging evidence indicating that KIF15 plays an important role in several malignancies including pancreatic cancer, hepatocellular carcinoma, and breast cancer (PMID: 33277366).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30774-1-AP (KIF15 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



WB result of KIF15 antibody (30774-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-KIF15 transfected HeLa cells.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using KIF15 antibody (30774-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).