

For Research Use Only

# HCN2 Polyclonal antibody

Catalog Number: 30946-1-AP



## Basic Information

Catalog Number:

30946-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG34038

GenBank Accession Number:

NM\_001194

GeneID (NCBI):

610

UNIPROT ID:

Q9UL51

Full Name:

hyperpolarization activated cyclic nucleotide-gated potassium channel 2

Calculated MW:

97 kDa

Observed MW:

60 kDa, 90 kDa, 120 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

## Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, rat, mouse

Positive Controls:

WB : mouse brain tissue, mouse cerebellum tissue, rat brain tissue

## Background Information

HCN2, also named as BCNG2, belongs to the potassium channel HCN family. The molecular weight of immature HCN2 is 90 kDa, and the protein is about 120 kDa after n-terminal glycosylation. It has been found that 60 kDa of HCN2 protein can be identified in mouse heart samples. Although all protein isolation steps were handled on ice and in the presence of protease inhibitors to reduce protease-dependent effects and to minimize protein degradation, it is certainly possible that the presence of the 60 kDa HCN2 protein in the heart samples does, in fact, reflect protein degradation. (PMID:21796099, PMID: 19574228)

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30946-1-AP (HCN2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.