For Research Use Only

## CD22 Polyclonal antibody

Catalog Number:30954-1-AP




Basic Information	Catalog Number: 30954-1-AP	GenBank Accession Number: NM_001771.4	Purification Method: Antigen affinity purification	
	Size: 500 µg/ml	GenelD (NCBI): 933	Recommended Dilutions: WB 1:2000-1:10000	
	Source: Rabbit	UNIPROT ID: P20273		
	lsotype: IgG	Full Name: CD22 molecule		
		Calculated MW: 95kd Observed MW: 120-130 kDa		
Applications	Tested Applications: WB, ELISA		Positive Controls: WB : Daudi cells, Raji cells, Ramos cells	
	Species Specificity: Human			
Background Informatio	is a 130-140 kDa, B-cell restr superfamily. The expression pro-B and pre-B cells and pre expression is lost during terr receptor for B-cell receptor (E	CD22, also known as Siglec-2 (sialic acid binding Ig-like lectin 2) or BL-CAM (B-lymphocyte cell adhesion molecule) is a 130-140 kDa, B-cell restricted, type I transmembrane glycoprotein belonging to the immunoglobulin gene superfamily. The expression of CD22 is developmentally regulated. It is expressed at low levels in the cytoplasm or pro-B and pre-B cells and present on the cell surface only at mature stages of B-cell differentiation. Cell surface expression is lost during terminal differentiation into plasma cell and after B-cell activation. CD22 is an inhibitory receptor for B-cell receptor (BCR) signalling, preferentially binds to alpha-2,6-linked sialic acid and mediates B-cell B-cell interactions. It plays a crucial role in activation and differentiation of the B-cell.		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage			

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30954-1-AP (CD22 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.