

## VCAM-1/CD106 Polyclonal antibody

Catalog Number: 30958-1-AP

2 Publications

## Basic Information

## Catalog Number:

30958-1-AP

## Concentration:

600 ug/ml

## Source:

Rabbit

## Isotype:

IgG

## GenBank Accession Number:

CAA47989.1

## GeneID (NCBI):

22329

## UNIPROT ID:

P29533

## Full Name:

vascular cell adhesion molecule 1

## Calculated MW:

81 kDa

## Observed MW:

100-110 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:2000-1:12000

IHC 1:200-1:800

IF-P 1:50-1:500

## Applications

## Tested Applications:

WB, IHC, IF-P, ELISA

## Cited Applications:

IHC, IF

## Species Specificity:

mouse, rat

## Cited Species:

mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

**WB** : rat spleen tissue, mouse spleen tissue, C2C12 cells

**IHC** : mouse spleen tissue, mouse liver tissue

**IF-P** : mouse spleen tissue,

## Background Information

Vascular cell adhesion molecule 1 (VCAM1), also known as CD106, is a 110-kDa transmembrane glycoprotein belonging to the immunoglobulin gene superfamily. VCAM1 is expressed by cytokine-activated endothelium, interacts with integrin VLA4 ( $\alpha 4 \beta 1$ ) present on the surface of leukocytes, and mediates both adhesion and signal transduction. It is also expressed either constitutively or inducibly in a variety of other cell types, including vascular smooth muscle cells, differentiating skeletal muscle cells, renal and neural epithelial cells, macrophages (Kupffer cells), dendritic cells, and bone marrow stromal cells (PMID: 7507076, 11359843).

## Notable Publications

Author	Pubmed ID	Journal	Application
Hui He	39721498	Redox Biol	IF
Zhuo Li	39490807	Microvasc Res	IHC

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30958-1-AP (VCAM-1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 30958-1-AP (VCAM-1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 30958-1-AP (VCAM-1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using VCAM-1 antibody (30958-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using VCAM-1 antibody (30958-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).