For Research Use Only

ADAMTS2 Polyclonal antibody

Catalog Number: 31032-1-AP



Basic Information

 Catalog Number:
 GenBank Accession Number:

 31032-1-AP
 NM_014244

 Size:
 GeneID (NCBI):

 400 ug/ml
 9509

 Source:
 UNIPROT ID:

 Rabbit
 095450

Isotype: Full Name:
IgG ADAM metallopeptidase with

Immunogen Catalog Number: thrombospondin
AG34957 Calculated MW:

135 kDa Observed MW: 110~130 kDa

thrombospondin type 1 motif, 2

Applications

Tested Applications: WB, IHC, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Positive Controls:

WB: A431 cells, A375 cells, PC-3 cells, THP-1 cell,

Purification Method:

WB 1:5000-1:50000 IHC 1:200-1:800

Antigen affinity Purification

Recommended Dilutions:

COLO 205 cells

IHC: mouse liver tissue,

Background Information

A disintegrin and metalloproteinase with thrombospondin motifs 2 (ADAMTS-2) is a metalloproteinase that plays a key role in the processing of fibrillar procollagen precursors into mature collagen molecules by excising the amino-propeptide. ADAMTS-2 can, like the procollagen C-proteinases, be regulated by transforming growth factor- β 1 (TGF- β 1), with implications for mechanisms whereby this growth factor effects net increases in formation of extracellular matrix. Mutations in the ADAMTS-2 gene cause dermatosparaxis in human (dermatosparactic type of Ehlers-Danlos syndrome, also known as Ehlers-Danlos syndrome type VIIC). Human ADAMTS-2 has eight potential sites for Asn-linked glycosylation and other mammals (PMID: 16046392, 12646579).

Storage


Storage:

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31032-1-AP (ADAMTS2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 31032-1-AP (ADAMTS2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 31032-1-AP (ADAMTS2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.