

For Research Use Only

DAB1 Polyclonal antibody

Catalog Number: 31459-1-AP



Basic Information

Catalog Number: 31459-1-AP	GenBank Accession Number: NM_021080	Purification Method: Antigen affinity Purification
Size: 500 ug/ml	GeneID (NCBI): 1600	Recommended Dilutions: WB 1:1000-1:4000
Source: Rabbit	UNIPROT ID: O75553	IF/ICC 1:200-1:800
Isotype: IgG	Full Name: disabled homolog 1 (Drosophila)	
Immunogen Catalog Number: AG35376	Calculated MW: 64 kDa	
	Observed MW: 45 kDa, 80 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls:
Species Specificity: human, pig	WB : pig cerebellum tissue, MCF-7 cells
	IF/ICC : HepG2 cells,

Background Information

Disabled-1 (Dab1) is an adaptor protein that is essential for neuronal migration and maturation in response to the extracellular protein Reelin. DAB1 protein docks to the intracellular part of the Reelin very low density lipoprotein receptor and apoE receptor type 2 and becomes tyrosine-phosphorylated following binding of Reelin to cortical neurons. The DAB1 protein contains a 180-amino acid N-terminal protein interaction/phosphotyrosine-binding (PTB) 1 domain that docks to the short cytoplasmic tail of the very low density lipoprotein receptor or apoE receptor type 2 at the level of NPXY motifs, with a preference for unphosphorylated motifs. Dab1 polypeptide bands ranging from 36 to 120 kDa have been identified in mouse embryonic brain, the 80-kDa Dab1 being the predominant form (PMID: 24313315).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31459-1-AP (DAB1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 31459-1-AP (DAB1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using DAB1 antibody (31459-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).