For Research Use Only

## NCAM1/CD56 Polyclonal antibody

Catalog Number: 31536-1-AP



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 31536-1-AP
 GeneID (NCBI):

 Size:
 17967

 350 µg/ml
 UNIPROT ID:

 Source:
 P13595-1

Source: P13595-1
Rabbit Full Name:

Isotype: neural cell adhesion molecule 1

G Calculated MW: 119 kDa

119 kDa Observed MW:

120 kDa, 140 kDa, 180 kDa

Purification Method: Antigen affinity Purification Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse brain tissue, mouse cerebellum tissue, rat

IHC: mouse brain tissue, mouse cerebellum tissue

## **Background Information**

Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronection-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isofroms have also been described (PMID: 1856291).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 31536-1-AP (NCAM1 antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 31536-1-AP (NCAM1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 31536-1-AP (NCAM1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).