For Research Use Only

CSDA Polyclonal antibody

Catalog Number: 31704-1-AP

Featured Product



Basic Information

Catalog Number: GenBank Accession Number: 31704-1-AP BC008801

Concentration: Genel D (NCBI):

 200 ug/ml
 8531

 Source:
 UNIPROT ID:

 Rabbit
 P16989

Isotype: Full Name:

IgG cold shock domain protein A

Immunogen Catalog Number: Calculated MW:

AG36519 40 kDa

Observed MW: 60-68 kDa Purification Method: Antigen affinity Purification Recommended Dilutions: WB 1:500-1:2000 IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IF/ICC, ELISA Species Specificity:

human, mouse

Positive Controls:

WB: mouse heart tissue, KYSE-30 cells, mouse liver

tissue, mouse skeletal muscle tissue

IF/ICC: U2OS cells,

Background Information

CSDA (Cold shock domain protein A), also known as DBPA (DNA-binding protein A), YBX3 (Y-box protein 3), is a DNA-binding protein that represses angiogenesis and lymphangiogenesis by directly binding to hypoxia response element (HRE) and serum response element (SRE). CSDA is ubiquitous expressed in human skeletal muscle, heart, and decidual cells, which is thought to a multifunctional epithelial-specific protein involved in gene transcription, DNA repair, and interaction with other proteins (PMID: 33816248). CSDA is a nucleic-acid-binding protein that directly regulates gene expression by using different mechanisms, and acts as a translational repressor (e.g., MYH9 and SPTBN1) and positive regulator of mRNA stability (PMID: 31189097). Western blot analysis detected CDSA at a range of 50-70 kDa due to the anomalous mobility of Y-box proteins (PMID: 38255791).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse heart tissue were subjected to SDS PAGE followed by western blot with 31704-1-AP (CSDA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of CSDA antibody (31704-1-AP; 1:1000; room temperature for 1.5 hours) with wild-type and CSDA knockout KYSE-30 cells.



Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using CSDA antibody (31704-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).