For Research Use Only

FRMD1 Polyclonal antibody Catalog Number: 31908-1-AP Featured Product

Featured Product



| Basic Information | Catalog Number: 31908-1-AP | GenBank Accession Number: BC137144 | Purification Method: Antigen affinity Purification |
|-----------------------|---|--|---|
| | Size: 500 ug/ml | GenelD (NCBI): 79981 | Recommended Dilutions: WB 1:1000-1:4000 |
| | Source: Rabbit | UNIPROT ID: Q8N878 | |
| | Isotype: IgG | Full Name: FERM domain containing 1 | |
| | Immunogen Catalog Number: AG36382 | Observed MW: 70 kDa | |
| Applications | Tested Applications: WB, ELISA | and the second | |
| | ND, ELISA Species Specificity: human, mouse, rat | | WB : mouse colon tissue, K-562 cells, mouse testis tissue, mouse thymus tissue, rat thymus tissue |
| Background Informatio | The FERM structural domain of FRMD1 (FERM Domain Containing 1) is widely found in cytoskeleton-associated proteins, which are capable of connecting the cytoskeleton to the cell membrane and is involved in signaling. FRMD1 is predicted to be involved in the positive regulation of the Hippo signaling pathway and may play a role in the cytoplasmic side of the cytoskeleton and apical plasma membrane. Although there are fewer studies on the role of FRMD1 in disease, its expression pattern in cancer and lymphoid tissues suggests that it may have important biological functions. | | |
| Storage | Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliguoting is unnecessary for -20° | I 50% glycerol pH 7.3. | |

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 31908-1-AP (FRMD1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. WB result of FRMD1 antibody (31908-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FRMD1 transfected K-562 cells.